

Singularity Free Inhomogeneous Models with Heat Flow

L. K. Patel* and Naresh Dadhich[†]**

*Department of Mathematics, Gujarat University,

Ahmedabad - 380 009, India

** Inter University Centre for Astronomy and Astrophysics,

Post Bag 4, Ganeshkhind, Pune - 411 007.

Abstract

We present a class of singularity free exact cosmological solutions of Einstein's equations describing a perfect fluid with heat flow. It is obtained as generalization of the Senovilla class [1] corresponding to incoherent radiation field. The spacetime is cylindrically symmetric and globally regular.

PACS numbers : 04.20 Jb, 98.80 Dr

[†] E-mail address : naresh@iucaa.ernet.in

Senovilla [1] has opened up a new vista of singularity free cosmological models. He obtained a class of cylindrically symmetric inhomogeneous exact solutions of Einstein's equations describing a perfect fluid with the equation of state $\rho = 3p$. The spacetime was globally regular, smooth and satisfied the energy and causality conditions. All the physical and curvature invariants were finite everywhere, and all causal curves were complete [2]. There was no singularity of any kind.

The singularity structure of a general class of inhomogeneous perfect fluid solutions was analysed [3] and it was shown that existence or non-existence of singularity as well as its kind (spacelike, timelike, Weyl, Ricci, etc) depended upon the particular choice of certain parameters and metric functions. There exists a general class of inhomogeneous perfect fluid solutions without singularity. Recently we have also found [4] a new class of solutions for stiff fluid ($\rho = p$) that have vacuum spacetime as the matter free ($\rho = 0$) limit. It is very curious that spacetime is empty as well as without singularity. One wonders about the source of its curvature.

How robust is the singularity free character of spacetime ? In particular what happens when dissipative effects of viscosity, heat flow etc. are incorporated ? In this note we wish to obtain a generalization of the Senovilla [1] model that includes heat flow.

Dissipative effects in cosmology were considered in the context of large entropy per baryon and isotropy of microwave background radiation[5,6] and heat flow was in particular investigated by several authors [7 - 16]. In very early times, matter is supposed to be highly dense and hot, and hence consideration of heat flow for very

early evaluation of the Universe is quite appropriate.

The energy momentum tensor incorporating heat flow with perfect fluid reads as

$$T_{ik} = (\rho + p)u_i u_k - p g_{ik} + 2u_{(i} q_{k)} \quad (1)$$

where $u_i u^i = 1$, $q_i u^i = 0$, and q_i is the heat flow vector. The expansion θ , shear scalar σ and the acceleration vector $\dot{u}_i$ for the fluid are defined by

$$\begin{aligned} \theta &= u^i_{;i}, \quad \sigma^2 = \sigma_{ik} \sigma^{ik} \\ \sigma_{ik} &= u_{(i;k)} - u_{(i} \dot{u}_{k)} - \frac{\theta}{3} (g_{ik} - u_i u_k). \\ \dot{u}_i &= u_{i;k} u^k \end{aligned}$$

For Einstein's equations

$$R_{ik} - \frac{1}{2} R g_{ik} = -8\pi T_{ik} \quad (2)$$

we obtain the following class of solutions for perfect fluid with radial heat flow;

$$ds^2 = A^2(dt^2 - dr^2) - B^2 dz^2 - D^2 d\phi^2 \quad (3)$$

where

$$\begin{aligned} A &= C^{1-b}(at) \quad C^{(b+4)/(1-2b)}(3ar) \\ B &= C^b(at) \quad C^{-(b+4)/9}(3ar) \\ D &= \frac{1}{3a} S(3ar) \quad C^{-(b+4)/9}(3ar) \quad C^{(1-b)}(at). \end{aligned} \quad (4)$$

Here a and b are free parameters and we denote $C(x) = \cosh x$, $S(x) = \sinh x$, $T(x) = \tanh x$, $Se(x) = \operatorname{sech} x$. The metric (3) is globally regular for entire

range of the coordinates; $-\infty < t, z < \infty, 0 < r < \infty, 0 < \phi < 2\pi$. The spacetime is cylindrically symmetric without any singularity.

The physical and kinematical parameters have the following explicit expressions:

$$8\pi\rho A^2 = \left(\frac{a^2}{3(1-2b)} \right) (b+4)(2b+17)(2-b)Se^2(3ar) + \frac{a^2}{3}(5-b)(b+1) + a^2(1-b^2)T^2(at) \quad (5.1)$$

$$8\pi p A^2 = \left(\frac{a^2}{9(1-2b)} \right) (b+4)^2(2b+17)Se^2(3ar) + \frac{a^2}{9}(b+1)(b+7) + a^2(1-b^2)T^2(at) \quad (5.2)$$

$$8\pi q_r A = \frac{4a^2}{3(1-2b)}(b+4)(2-b)(b+1)T(at)T(3ar) \quad (5.3)$$

$$\theta A = a(2-b)T(at) \quad (5.4)$$

$$\sigma^2 A^2 = \frac{2}{3}a^2(1-2b)^2T^2(at) \quad (5.5)$$

$$\dot{u}_r = -\frac{3a}{1-2b}(b+4)T(3ar). \quad (5.6)$$

The phenomenological expression for the heat conduction is given by

$$q_k = \psi(F_{,i} + F\dot{u}_i)(\delta_k^i - u^i u_k) \quad (6)$$

where ψ is the thermal conductivity and F is the temperature. For the case under consideration, the above equation can be integrated if ψ is a function of t alone.

The temperature distribution in the Universe is then given by

$$F = l_0(t)C^{(b+4)/(1-2b)}(3ar) - \frac{a}{36\pi\psi}(2-b)(b+1)T(at) C^{-(b+4)/(1-2b)}(3ar) \quad (7)$$

where l_0 is an arbitrary function of t .

The solution(3) reduces to the Senovilla [1] perfect fluid model for $b = -1$, when heat flux is switched off ($q_k = \psi = 0$). From the expressions in equation (5) the physical requirements ($\rho > p > 0$, $\rho + 3p > 0$) restrict the range for b to $-1 \leq b < 1/2$. All the parameters remain finite and well behaved for all values of t and r . We now compute the Weyl scalars and they are as follows :

$$\begin{aligned} \psi_0 + \psi_4 = & C^{2(b-1)}_{(at)} C^{\frac{2(b+4)}{2b-1}}_{(3ar)} \left[\frac{a^2}{(1-2b)}(39 + 5b - 4b^2) \right. \\ & \left. + (1-b)(1-2b)a^2T^2(at) \right] \end{aligned}$$

$$\psi_0 - \psi_4 = \frac{2}{3}a^2(5-b)(b+4)T(at)T(3ar)C^{2(b-1)}_{(at)}C^{\frac{2(b+4)}{2b-1}}_{(3ar)}$$

$$\begin{aligned} \psi_2 = & \frac{1}{6}C^{2(b-1)}_{(at)} C^{\frac{2(b+4)}{2b-1}}_{(3ar)} \left[\frac{a^2}{1-2b}(35 + 13b - 4b^2) \right. \\ & \left. + (1-3b+2b^2)a^2T^2(at) - \frac{2(20+b-b^2)}{(1-2b)}a^2T^2(3ar) \right] \end{aligned}$$

It is clear from the above expressions that the Weyl scalars remain regular and finite for the entire range of coordinates for $-1 \leq b < \frac{1}{2}$.

From relations (5) it follows that as $t \rightarrow -\infty$ all the physical and kinematic parameters, as well as the Weyl curvatures tend to zero, however the metric does not go over to flat spacetime (this also happens in [1]). Then as t increases, the fluid starts contracting, the density and pressure increase reaching the maximum at

$t = 0$. The heat flux q_r and expansion θ are negative for $-\infty < t < 0$, zero at $t = 0$ and positive for $0 < t < \infty$. Similarly the shear is positive in r and ϕ directions and negative in z direction and it reverses its sense at $t = 0$. The radial acceleration is zero at $r = 0$ and negative for $r > 0$. As t increases from zero, the fluid starts expanding and heat flows radially out. All physical and kinematical parameters tend to zero again as $t \rightarrow \infty$ and the spacetime tends to the same state as that of $t \rightarrow -\infty$. The universe therefore starts from nearly flat spacetime at $t \rightarrow -\infty$, passes through the dense and hot state at $t = 0$ (the free parameter a can be chosen as large as we please) and becoming nearly flat and cold again as $t \rightarrow \infty$.

We have thus shown that it is possible to incorporate heat flux in the Senovilla class [1] of models without disturbing their singularity free character. We have also subsequently generalised the general class of perfect fluid solution [3] to include heat flux [17].

References

- [1] Senovilla J.M.M., 1990, Phys. Rev. Lett. *64*, 2219.
- [2] Chinea, F.J., Fernandez-Jambrina, L. and Senovilla, J.M.M., 1992, Phys. Rev., *D45*, 481.
- [3] Ruiz, E. and Senovilla, J.M.M., 1992, Phys. Rev. *D45*, 1995.
- [4] Patel L.K. and Dadhich N., 1992 ,Preprint IUCAA-1/93 Jan'93
- [5] Misner, C.W., 1968, Ap. J. *151*, 431.
- [6] Caderni, N. and Fabri, R., 1978, Nuovo Cimento, *44B*, 228.
- [7] Dang, Y., 1989, Gen. Rel. Grav., *21*, 503.
- [8] Mukherjee, G., 1986, J. Astrophys. Astron., *7*, 259.
- [9] Novello, M. and Reboucas, M.J., 1978, Ap. J., *225*, 719.
- [10] Ray, D., 1980, J. Math. Phys., *21*, 2797.
- [11] Reboucas, M.J. and de Limma, J.A.S., 1982, J. Math. Phys. *23*, 1547.
- [12] Reboucas, M.J. and de Limma, J.A.S., 1981, J. Math. Phys. *22*, 2699.
- [13] Reboucas, M.J., 1980, Nuovo Cimento, *67B*, 120.
- [14] Bradley, J.M. and Sveistins, E., 1984, Gen. Rel. Grav. *16*, 1119.
- [15] Sveistins, E., 1985, Gen. Rel. Grav., *17*, 521.
- [16] Patel, L.K. and Dadhich, N., 1991, Ap.J. *401*, 433.
- [17] Patel, L.K. and Dadhich, N., to be published.